

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 August 2003 (28.08.2003)

PCT

(10) International Publication Number  
**WO 03/070246 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 31/496**,  
A61P 25/00
- (21) International Application Number: PCT/IB03/00642
- (22) International Filing Date: 17 February 2003 (17.02.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/358,806 20 February 2002 (20.02.2002) US  
60/359,038 21 February 2002 (21.02.2002) US  
60/360,459 27 February 2002 (27.02.2002) US
- (71) Applicant: **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).
- (72) Inventors: **BUSCH, Frank, Robert**; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **GROBIN, Adam, Worth**; 109 Cardiff Place, Chappel Hill, NC 27516 (US). **HOWARD, Harry, Ralph, Jr.**; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **LEEMAN, Kyle, Robert**; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US).
- (74) Agents: **LUMB, J., Trevor et al.**; Pfizer Inc., 201 Tabor Road, Morris Plain, NJ 07950 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: CONTROLLED SYNTHESIS OF ZIPRASIDONE AND COMPOSITIONS THEREOF

(57) Abstract: The subject invention provides a ziprasidone composition that comprises not greater than 1000 ppm des-chloro ziprasidone, preferably not greater than about 500 ppm des-chloro ziprasidone, and more preferably not greater than about 100 ppm des-chloro ziprasidone. Methods for synthesizing and using such ziprasidone compositions are also provided.

WO 03/070246 A1

## CONTROLLED SYNTHESIS OF ZIPRASIDONE AND COMPOSITIONS THEREOF

Background of the Invention

Efforts are made to prepare pharmaceutical products of a high grade and with a minimum amount of impurities present. The control of impurities requires a study of various options to decide upon the reaction conditions and testing protocols necessary to insure that drugs which are administered to the public are pure.


Guidance given by regulatory bodies, including the United States Food and Drug Administration (FDA), suggests that impurities in drugs be identified, if present, if they are at a level of 0.1% (that is, 1000 ppm) or greater for drug substances dosed at 2g/day or lower. (Note that ppm is parts per million, so that 1% = 10,000 ppm; 0.1% = 1000 ppm; 0.01% = 100 ppm; and 0.001% = 10 ppm). For example, the FDA has indicated that identification of impurities below apparent levels of 0.1% for a 2g/day-dosed drug substance is generally not considered necessary (Federal Register Vol. 65, No. 140 pp. 45085-45090, 45086 and 45089 (July 20, 2000)). However, the FDA also points out that tighter controls may be necessary for some impurities, depending upon their specific properties (*Id.* at 45086). Furthermore, studies to obtain safety information for a proposed quantity of an impurity are recommended if the proposed quantity exceeds a qualification threshold of 0.05% (500 ppm for a drug substance dosed at 2g/day or lower (*Id.* at 45087 and 45089)).

Ziprasidone (5-(2-(4-(1,2-benzisothiazol-3-yl-1-piperazinyl)-ethyl)-6-chloro-1,3-dihydro-2-(1H)-indol-2-one) is a potent antipsychotic agent and is useful for treating various disorders including schizophrenia, anxiety and migraine pain. Ziprasidone has been approved by the FDA for treatment of schizophrenia and goes by the brand name Geodon<sup>TM</sup> in the United States. Ziprasidone has also been indicated as useful for treating Tourette's Syndrome (United States Patent 6,127,373), glaucoma and ischemic retinopathy (EP 985414 A2), and psychiatric conditions including dementia of the Alzheimer's type, bipolar disorders, mood disorders, panic disorders, agoraphobia, social phobia, panic disorder, post-traumatic stress disorder, acute stress disorder, substance-induced anxiety disorder, anxiety disorders not otherwise specified, dyskinesias and behavioral manifestations of mental retardation, conduct disorder, and autistic disorder (United States Patent 6,245,766). The aforementioned United States Patents and European Patent Application are all incorporated by reference herein in their entireties.

United States Patent 4,831,031 describes a genus of compounds encompassing ziprasidone and the synthesis of such compounds. Another method for synthesizing ziprasidone is described in United States Patent 5,206,366. A method for specifically synthesizing ziprasidone hydrochloride monohydrate is

described in United States Patent 5,312,925. A method for synthesizing ziprasidone mesylate dihydrate is described in United States Patent 6,245,765; and a method for synthesizing ziprasidone mesylate trihydrate is described in United States Patent 6,110,918. United States Patents 5,338,846; 5,359,068; and 6,111,105 also describe methods for synthesizing ziprasidone and/or intermediates therefore. The  
5      aforementioned United States Patents are all incorporated by reference herein in their entireties.

The structure of ziprasidone can be depicted as:



10      (H. Howard, et al., "Ziprasidone Hydrochloride", Drugs of the Future 1994, 19(6): 560-563. As can be seen from the structure above, the compound ziprasidone comprises a chlorine atom.

Methods of introducing halogens into organic compounds are summarized in many organic text books. For example, J. March, Advanced Organic Chemistry, 4<sup>th</sup>  
15      Edition, pp. 587-591, and references cited therein, has a discussion of halogenation chemistry. More specifically, formation of chloro-aromatic compounds are frequently formed by a variety of methods also well known to those skilled in the art, and again summarized in J. March, Advanced Organic Chemistry, 4<sup>th</sup> Edition, Chapter 11, "Aromatic Electrophilic Substitution". The chemistry to add a halogen, or more  
20      specifically a chlorine, to an aromatic group is thus well known to those skilled in the art. It is also known that such chemistry usually results in some mixtures of molecules, one of which is commonly the unreacted starting material not containing the chlorine atom. Further, over-chlorination is a problem well known to those skilled in the art; it is common to form some dichloro-compound impurities when the mono-chloro is desired and some trichloro-compound impurities when the dichloro- is  
25      desired. Over-chlorination is typically controlled by limiting the amount of the chlorinating reagent used. Unfortunately, control of over-chlorinated analogs in the drug substance by limiting the amount of chlorinating reagent utilized in the

introduction of the aromatic chlorine substituent would be expected to result in more of a des-chloro impurity (unreacted starting material not containing the chlorine atom).

#### Summary of the Invention

5 The des-chloro analog of ziprasidone is 5-[2-[4-(1,2)-benzothiazol-3-yl]-1-piperazinyl]ethyl]-1,3-dihydro-2H-indol-2-one (hereinafter des-chloro ziprasidone). Based on the known methods for synthesizing halogenated aromatic compounds referred to above, any synthesized batch of ziprasidone drug substance will comprise some amount of a des-chloro ziprasidone impurity. Control of over-chlorinated  
10 analogs in the drug substance by limiting the amount of chlorinating reagent utilized in the introduction of the aromatic chlorine substituent would be expected to result in more of the des-chloro ziprasidone impurity.

The subject invention pertains to the techniques we have developed to control the synthesis of ziprasidone drug substance to insure that levels of des-chloro ziprasidone are at low levels. In our particular drug substance for use in  
15 pharmaceutical compositions, a level of not greater than about 100 ppm des-chloro ziprasidone consistently is met. However, our invention pertains to ziprasidone compositions comprising levels of des-chloro ziprasidone of up to but not greater than about 1000 ppm and to methods for controlling the levels of des-chloro ziprasidone to up to but not greater than about 1000 ppm in a ziprasidone composition.

20 This invention further relates to a ziprasidone composition comprising low levels of des-chloro ziprasidone, preferably not greater than about 1000 ppm des-chloro ziprasidone, more preferably not greater than about 500 ppm des-chloro ziprasidone, and even more preferably not greater than about 100 ppm des-chloro ziprasidone.

25 As used herein and unless otherwise indicated, the term "ziprasidone" includes ziprasidone free base and pharmaceutically acceptable salts of ziprasidone. A generic teaching of preparation of pharmaceutically acceptable salts of a genus of compounds including ziprasidone is provided in United States Patent 4,831,031 (see, for example, Column 3 thereof), incorporated herein by reference. In one  
30 embodiment, the ziprasidone in the composition of the present invention is ziprasidone free base. In another embodiment, the ziprasidone in the composition of the present invention is ziprasidone hydrochloride monohydrate. In another embodiment, the ziprasidone is ziprasidone mesylate dihydrate, and in another embodiment, the ziprasidone is ziprasidone mesylate trihydrate.

The term "ziprasidone drug substance", as used herein and unless otherwise indicated, refers to a ziprasidone composition, as defined above, that is used in the formulation of a pharmaceutical composition. Such pharmaceutical composition may contain pharmaceutical carriers, excipients, flavorings, and other ingredients that are known to be used in pharmaceutical compositions and as described in more detail below.

This invention also provides a pharmaceutical composition for treating in a mammal, including a human, a disorder or condition selected from schizophrenia, anxiety, migraine pain, Tourette's Syndrome, glaucoma, ischemic retinopathy, dementia of the Alzheimer's type, a bipolar disorder, a mood disorder, agoraphobia, social phobia, panic disorder, post-traumatic stress disorder, acute stress disorder, substance-induced anxiety disorder, an anxiety disorders not otherwise specified (NOS), dyskinesias, a behavioral manifestation of mental retardation, conduct disorder, and autistic disorder comprising an amount of a ziprasidone drug substance which is a composition comprising ziprasidone and an amount of des-chloro-ziprasidone of not greater than about 1000 ppm, which amount of ziprasidone drug substance is effective in treating said disorder or condition, and a pharmaceutically acceptable carrier.

In one embodiment, the amount of des-chloro ziprasidone in the ziprasidone drug substance is not greater than about 500 ppm. In a preferred embodiment, the amount of des-chloro ziprasidone in the ziprasidone drug substance is not greater than about 100 ppm des-chloro ziprasidone.

This invention also provides a method for treating in a mammal, including a human, in need thereof a disorder or condition selected from schizophrenia, anxiety, migraine pain, Tourette's Syndrome, glaucoma, ischemic retinopathy, dementia of the Alzheimer's type, a bipolar disorder, a mood disorder, agoraphobia, social phobia, panic disorder, post-traumatic stress disorder, acute stress disorder, substance-induced anxiety disorder, an anxiety disorders not otherwise specified (NOS), dyskinesias, a behavioral manifestation of mental retardation, conduct disorder, and autistic disorder, which method comprises administering to said mammal an amount of a ziprasidone drug substance which is a composition comprising ziprasidone and an amount of des chloro ziprasidone of not greater than about 1000 ppm, which amount of ziprasidone drug substance is effective in treating said disorder or condition.

In one embodiment, the amount of des-chloro ziprasidone in the ziprasidone drug substance is not greater than about 500 ppm. In a preferred embodiment, the amount of des-chloro ziprasidone in the ziprasidone drug substance is not greater than about 100 ppm des-chloro ziprasidone.

5 This invention also provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises starting with a composition of a chlorinated reactant comprising a sufficiently low level of non-chlorinated impurity to synthesize said ziprasidone composition. In one embodiment, the chlorinated  
10 reactant is a composition of 6-chlorooxindole (6-chloro-1,3-dihydro-2H-indol-2-one).

In a more specific embodiment, this invention provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

15 a) obtaining one or more samples of one or more 6-chloro-1,3-dihydro-2H-indol-2-one batches;

b) measuring the level of oxindole impurity in each of the samples of (a);

c) selecting a 6-chloro-1,3-dihydro-2H-indol-2-one batch that comprises a level of oxindole of not greater than about 0.3% based on the measurement or measurements conducted in (b); and

20 d) using the batch selected in (c) to synthesize said ziprasidone composition.

In one embodiment, step (c) comprises selecting a 6-chloro-1,3-dihydro-2H-indol-2-one batch that comprises a level of oxindole of not greater than about 0.15%. In a preferred embodiment, step (c) comprises selecting a 6-chloro-1,3-dihydro-2H-indol-2-one batch that comprises a level of oxindole of not greater than about 0.03%.

25 Although there are many known routes to 6-chlorooxindole, starting materials therefore are typically a substituted 4-chlorotoluene or 1,4-dichloro-nitrobenzene (see, G. J. Quallich and P. M. Morrissey, Synthesis, 1993, 51-53; and references cited therein; and F. R. Busch and R. J. Shine, "Development of an Efficient Process to 6-Chlorooxindole", presented at the 208th ACS National Meeting in Washington D.C. in the Symposium on Technical Achievements in Organic Chemistry, 1994, (talk #126).). However, the concept of controlling chlorinated isomers, over-chlorination, or des-chloro impurities for the synthesis of 6-chlorooxindole is not described in the prior art. G. J. Quallich and P. M. Morrissey, *supra*, is incorporated herein by  
30 reference in its entirety. Other methods of synthesizing 6-chlorooxindole can be



determined by a person of ordinary skill in the art, and such methods are included in the step of obtaining a batch of 6-chlorooxindole for the above-described method of this invention. Furthermore, a batch of 6-chlorooxindole can be obtained by purchase from manufacturers of organic chemicals, for example Plaistow, Ltd., Little Island, County Cork, Ireland or Finorga, Route de Givors, 38670 Chasse-Sur-Rhone, France.

As used herein, a "6-chlorooxindole batch" is a composition consisting essentially of 6-chlorooxindole, which composition may contain low levels of impurities, one of which may be oxindole.

The level of oxindole impurity in a sample of a batch of 6-chlorooxindole can be determined using standard analytical techniques known to those of ordinary skill in the art. For example, the level of oxindole impurity may be determined by normal phase HPLC, reverse phase HPLC, or gas chromatography methods.

A specific method for determining the level of oxindole in a sample of a 6-chlorooxindole batch is referred to herein as "Detection Method B" and is provided in the "Detailed Description of the Invention" Section of this application below.

This invention also provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

a) acylating a composition comprising 6-chloro-1,3-dihydro-2H-indol-2-one and an oxindole impurity with chloroacetyl chloride by Friedel-Crafts Acylation to synthesize a composition comprising 6-chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one;

b) treating the composition resulting from (a) to reduce of the oxo of the chloroacetyl group therein to form a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity;

c) isolating a sample of the composition resulting from (b);

d) measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity in the isolated sample from (c);

e) determining whether or not the quantity in (d) is not greater than about 0.28%; and

f) purifying by recrystallization and/or reslurry the composition resulting from (b) if the quantity measured in (d) is greater than about 0.28% until the quantity of the

5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity is not greater than about 0.28%, and synthesizing a ziprasidone composition from the composition so purified; or,

g) if the quantity in (d) is not greater than about 0.28%, synthesizing a ziprasidone composition from the composition of (b).

5 In a preferred embodiment, the composition of ziprasidone prepared according to the method in the preceding paragraph comprises an amount of des-chloro ziprasidone of not greater than about 500 ppm, with the value of 0.28% provided in steps (e) and (f) being adjusted accordingly to about 0.14%. In a more preferred embodiment, the composition of ziprasidone prepared according to the  
10 method in the preceding paragraph comprises an amount of des-chloro ziprasidone of not greater than 100 ppm, with the value of 0.28% in steps (e) and (f) being adjusted accordingly to about 0.028%.

Measurement of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one, as in step (d) in the method described in the preceding paragraphs, can be conducted by standard  
15 analytical chemistry techniques, for example reverse phase HPLC or other suitable chromatographic methods.

However, in a preferred embodiment, the 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one is measured in step (b) by Detection Method A, described below.

In one embodiment, the purification of a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in step (f) is by reslurry. Reslurry is a  
20 process similar to recrystallization, but where all of the material is not completely dissolved. The composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in step (f) may however be purified by recrystallization, reslurry, or a combination thereof. A preferred method of purification of 6-chloro-5-(2-chloroethyl)-  
25 1,3-dihydro-2H-indol-2-one is by recrystallization and/or reslurry in an aqueous miscible solvent, preferably acetonitrile/water.

This invention also provides a HPLC method, called "Detection Method A" herein, for measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one.

30 More specifically, this invention provides a method, "Detection Method A", using HPLC for measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one, which method comprises

a) preparing sample solution from said composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one by dissolving a portion of said composition in an organic solvent, followed by dilution with an organic solvent of the dissolved portion so that a concentration (weight/volume), based on the weight of said portion and volume of solvent, of about 1 mg/mL is obtained;

b) running the sample solution through a stable-bond cyano HPLC column using a mobile phase consisting essentially of (75:13-17:8-12 v/v/v/v) 0.05 M  $\text{KH}_2\text{PO}_4$ , pH=from 5.5-6.5,:Acetonitrile:Methanol; at a column temperature of from 30° C to 40° C; with detection by UV light at 254nm UV;

c) detecting a peak appearing at from between 8 to 10 minutes on a chromatogram resulting from (b);

d) measuring the peak area (named  $A_c$ ) of the peak detected in (c);

e) preparing a standard from a composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one by dissolving and diluting a portion of said composition in an organic solvent such that the concentration (weight/volume) of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one, based on the weight of the portion and volume of the solvent, is about equal to a selected fraction value at or above which detection of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is desired;

f) running the standard through a stable-bond cyano HPLC column using a mobile phase consisting essentially of (75:13-17:8-12 v/v/v/v) 0.05 M  $\text{KH}_2\text{PO}_4$ , pH=from 5.5-6.5,:Acetonitrile:Methanol; at a column temperature of from 30° C to 40° C; with detection by UV light at 254nm UV;

g) measuring the peak area (named  $A_{\text{pur1}}$ ) of the peak on a chromatogram resulting from (f); and

h) calculating the quantity of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one by

i) calculating the Response Factor for 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one according to the following formula:

$$R_{\text{pur1}} = (A_{\text{pur1}})(DF)/(W_{\text{pur1}})(PF)$$

wherein :  $A_{\text{pur1}}$  is as defined above;

$W_{\text{pur1}}$  = weight of composition in the standard;

PF = potency factor for 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one; and

DF = dilution factor for the standard; and

ii) calculating the %w/w of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one according to the following formula:

$$\%w/w = (A_c)(DF)(100)/(R_{pur1})(W_{S2})$$

wherein :  $A_c$  is as defined above;

$R_{pur1}$  = Response Factor calculated in (h)(i) above;

$W_{S2}$  = weight of the portion of the composition used in step (a); and

DF = dilution factor for sample solution.

Organic solvents that are useful in steps (a) and (e) include, but are not limited to THF (tetrahydrofuran), methanol, acetonitrile, or the mobile phase described in step (a). Other organic solvents may also be useful in this method. In a preferred embodiment of the HPLC method described in the preceding paragraph, the sample solution is dissolved in THF and is subsequently diluted using the mobile phase. In another preferred embodiment, the standard is prepared by diluting the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in THF.

In another preferred embodiment, a flow rate of about 1.0mL/min is used in the HPLC method. In another preferred embodiment, injection volumes of the standard and of the sample solution of at least about 20  $\mu$ L, more preferably about 20  $\mu$ L, are used.

In another preferred embodiment of the above-described HPLC method, the column temperature is 35° C. In another preferred embodiment, the ratio of  $\text{KH}_2\text{PO}_4$ :acetonitril:methanol is 75:15:10. In another preferred embodiment, the pH of the  $\text{KH}_2\text{PO}_4$  is 6.0.

A "stable-bond cyano column", as used herein, means a HPLC column comprising a stationary phase consisting essentially of a cyano bonded phase. Stable-bond cyano columns are known to those of ordinary skill in the art. Such HPLC columns are readily available to those of ordinary skill in the art from

commercial sources, for example the HPLC column Zorbax™ (Mac-Mod Analytical, P.O. Box 2600, 127 Commons Court, Pennsylvania 19317, USA).

5 The "Potency Factor" used in the calculation in step (h) in the above-described method refers to the purity of the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one with respect to the 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one therein. The Potency Factor can be determined by a person of ordinary skill in the art by deducting quantities, if any, of substances for which detection is typically conducted by a person of ordinary skill when analyzing the purity of an organic composition. Such substances include, for example, water, solvent or solvents, and "residue on ignition" (i.e. inorganic matter, for example sodium or potassium). Detection and quantification of such substance can be determined by a person of ordinary skill in the art. Hence, taking into account such substances, a Potency Factor for a composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one may be, for example, 98% 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one.

10 The "Dilution Factors" (for the sample solution and for the standard) used in the calculation in step (h) in the above-described method refer to the amount by which the composition being analyzed and the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one was diluted in steps (a) and (e), respectively. Hence, the Dilution Factor for the sample solution will be the volume of solvent used to prepare the sample solution in step (a) of the method. For example, if 80 mg of a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is used in step (a), and 80 mL of solvent accordingly, then the Dilution Factor will be 80. The Dilution Factor for Standard A will depend on the value selected in step (e). For example, if detection of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one at or above about 100 ppm is selected, then the concentration of the 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the standard will be about 0.0001. If, for example, 20 mg of the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is used in step (e), then the Dilution Factor for the standard is 20/0.0001, or  $2 \times 10^5$ . As another example, if detection of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one at or above about 500 ppm is selected, then the concentration of the 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the standard will be about 0.0005. If, for example, 20 mg of the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is used in step (e), then the Dilution Factor for the standard will be 20/0.0005, or  $4 \times 10^4$ .

As another example, if detection of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one at or above about 1000 ppm is selected, then the concentration of the 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the standard will be about 0.001. If, for example, 20 mg of the composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is again used in step (e), then the Dilution Factor for the standard will be 20/0.001, and the Dilution Factor for the standard will be  $2 \times 10^4$ .

This invention also provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

a) reducing a composition comprising 6-chloro-5-(chloroacetyl)-1,3-dihydro-2*H*-indol-2-one and a 5-(chloroacetyl)-1,3-dihydro-2*H*-indol-2-one impurity by treatment with triethylsilane in the presence of a strong acid to obtain a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity; and

b) synthesizing a composition comprising ziprasidone from the composition resulting from (a). In a preferred embodiment, the strong acid in step (a) comprises trifluoroacetic acid or methanesulfonic acid.

In another preferred embodiment, the method further comprises

i) isolating prior to step (b) a sample of the composition from (a), and measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity in said sample;

ii) determining whether or not the quantity in (i) is not greater than about 0.28%; and

iii) purifying by recrystallization and/or reslurry the composition from (a) if the quantity in (i) is greater than about 0.28% until the quantity of the 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity is not greater than about 0.28%, and then proceeding to step (b) using the composition from (a) so purified; or

iv) if the quantity in (i) is not greater than about 0.28%, then proceeding to step (b).

In another embodiment, the method is for synthesizing a ziprasidone composition comprising not greater than 500ppm of des-chloro ziprasidone. The value of 0.28% used for analyzing the sample from step (a) may be adjusted accordingly to 0.14%. In another embodiment, the aforementioned method is for synthesizing a ziprasidone composition comprising not greater than 100ppm. The

value of 0.28% used for analyzing the method may likewise be adjusted accordingly to 280 ppm.

In another embodiment of this method, the quantity of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in the sample of the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one is measured by a method comprising Detection Method A.

The purification of the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one in step (a)(ii) of the above method is by reslurry and/or recrystallization. The recrystallization and/or reslurry are in a suitable solvent mixture, preferably an aqueous miscible solvent, and more preferably a mixture of acetonitrile/water.

This invention also provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

a) purifying a composition comprising 6-chloro-1,3-dihydro-2*H*-indol-2-one and an oxindole impurity until a composition comprising not greater than about 0.3% of said oxindole impurity is obtained; and

b) using the composition resulting from (a) to synthesize a ziprasidone composition.

In one embodiment, the composition in (a) is purified until a composition comprising not greater than about 0.15% of oxindole impurity is obtained, and a composition that comprises an amount of des-chloro ziprasidone of not greater than about 500 ppm is synthesized.

In another embodiment, the composition in (a) is purified until a composition comprising not greater than about 0.03% of oxindole impurity is obtained, and a composition that comprises an amount of des-chloro ziprasidone of not greater than about 100 ppm is synthesized.

Methods of purification of the composition comprising -chloro-1,3-dihydro-2*H*-indol-2-one and an oxindole impurity that can be used in this invention include extraction and/or recrystallization and/or reslurry. In one embodiment the purification is by recrystallization and/or reslurry from organic solvents. See, for example, G. J. Quallich and P. M. Morrissey, *supra*, and references cited therein; and F. R. Busch and R. J. Shine, *supra*.

This invention also provides a method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

5 a) recrystallizing and/or reslurrying a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity until a composition comprising not greater than about 0.3% of said 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity is obtained; and

b) using the composition resulting from (a) to synthesize a ziprasidone composition.

10 In one embodiment, the composition in (a) is recrystallized and/or reslurried until a composition comprising not greater than about 0.15% of said 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity is obtained, and a composition that comprises an amount of des-chloro ziprasidone of not greater than about 500 ppm is synthesized.

15 In another embodiment, the composition in (a) is recrystallized and/or reslurried until a composition comprising not greater than about 0.03% of 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity is obtained, and a composition that comprises an amount of des-chloro ziprasidone of not greater than about 100 ppm is synthesized.

20 In a preferred embodiment, the recrystallization/reslurrying conditions in step (a) of this method comprise use of polar organic solvents and/or polar organic solvents mixed with a small volume of water. Preferably, the recrystallization and/or reslurry is with an aqueous miscible solvent, for example acetonitrile/water.

25 The terms "treatment", "treating", and the like, refers to reversing, alleviating, or inhibiting the progress of the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. As used herein, these terms also encompass, depending on the condition of the patient, preventing the onset of a disorder or condition, or of symptoms associated with a disorder or condition, including reducing the severity of a disorder or condition or symptoms associated therewith prior to affliction with said disorder or condition. Thus, "treatment", as used herein, can refer to administration of a compound of the invention to a subject that is not at the time of  
30 administration afflicted with the disorder or condition. "Treating" thus also encompasses preventing the recurrence of a disorder or condition or of symptoms associated therewith.



"Mammal", as used herein, and unless otherwise indicated, means any mammal. The term "mammal" includes, for example and without limitation, dogs, cats, and humans.


5 The term "about", when used herein in, for example, "less than 'about' 1000 ppm" means within a range of plus or minus 10% of the value to which the term is being applied.

Detailed Description of the Invention


10 One method for synthesizing ziprasidone, as noted above, is taught in United States Patent 5,206,366, which has been incorporated herein by reference. This Patent teaches that ziprasidone can be synthesized according to the method depicted in Scheme 1, below. Scheme 2, below, illustrates the mechanism by which des-chloro ziprasidone forms from any oxindole impurity in the starting reactant if a synthesis according to Scheme 1 is conducted. "IPO" in the following Schemes indicates isopropyl alcohol:

Scheme 1

Step 1




Step 2




5


Step 3



16



5

Scheme 2

10

One control strategy we identified was determining the purging of the des-chloro impurities during chemical synthesis of ziprasidone, and then setting sufficient limits on the quality of the starting reactant used to synthesize ziprasidone. During a synthesis of ziprasidone, one or more of the intermediates undergoes extraction and crystallization during the reaction work-ups. Although each intermediate and its corresponding des-chloro impurity are structurally similar and have similar solubility, there are slight differences in the physio-chemical properties. Thus, we conducted

15

purging experiments to determine the amount of des-chloro impurity that is removed by the extraction and recrystallization processing operations.

Referring to Schemes 1 and 2, Compound 6, oxindole, a potential impurity in a batch of the starting material, Compound 1, 6-chlorooxindole, may proceed under reaction conditions to result in compound 9, des-chloro ziprasidone. Surprisingly, we found a greater than three-fold purge of Compound 6 and its subsequent analogs during the reaction and isolation conditions of the synthetic process. Therefore, commercial supplies of Compound 1 containing up to 0.03% of Compound 6 will result in ziprasidone drug substance containing less than 0.01% (100 ppm) of Compound 9.

The control of a maximum limit of about 0.03% (300 ppm) Compound 6 in the purchased Compound 1 is justified based on the levels of Compound 6 detected in previous lots of Compound 1, purging information obtained during the development of the synthesis, and measurements obtained during the processing of these intermediates.

As noted above, oxindole can be detected by standard analytical techniques. One specific method we have found for determining oxindole in a composition comprising 6-chlorooxindole is as follows and will be referred to herein as "Detection Method B":

#### DETECTION METHOD B:

##### PRINCIPLE:

Normal phase liquid chromatography (LC) is used to separate Compound 1 from its potential impurities. Comparison of the peak areas and retention times for samples and the working standard of Compound 1 provides a quantitative assay and identification test for Compound 1. Comparison of the peak areas of the specified impurities, if present, against dilute solutions of the impurities, provides a quantitative measure of their abundance.

##### APPARATUS:

1. Standard laboratory equipment.
2. Suitable liquid chromatograph
  - a. Pump - constant flow delivery
  - b. UV detector - 254 nm
  - c. Injector capable of making 50 µL injections

d. Data acquisition system

3. Column:

Waters Associates Nova-Pak silica column, 4 micron particles, 150 x

3.9 mm

5

(i.d.).

(2 columns placed in series or equivalent length of 300 mm), or

equivalent

4. Balance capable of weighing 50 mg  $\pm$  0.1 mg - e.g. Mettler AE240

**REAGENTS:**

10

1. Hexane - HPLC grade

2. Tetrahydrofuran (THF)

3. Isopropanol (IPO) - HPLC grade

4. 15-crown-5 (1,4,7,10,13-pentaoxacyclopentadecane)

5. Working standard of **Compound 6** (oxindole)

15

**CHROMATOGRAPHIC CONDITIONS:**

1. Flow Rate: 1.5 mL/min

2. Injection Volume: 50  $\mu$ L

3. Detection Wavelength: 254 nm

4. Mobile Phase: hexane / isopropanol / tetrahydrofuran / 15-crown-5

20

1000 / 9 / 9 / 0.5 (v/v/v/v)

5. Column Temperature: Ambient (about 23°C)

Under the above cited conditions, **Compound 6** will elute between 15 - 19 minutes.

The elution time for **Compound 1** is between 17 - 21 minutes.

**MOBILE PHASE PREPARATION:**

25

In a 2 liter flask, add in the following order: 1000 mL of hexane, 9 mL of isopropanol, 9 mL of tetrahydrofuran, and 0.5 mL of 15-crown-5. Mix well and degas under vacuum with sonication or stirring for about 20 seconds.

**SAMPLE AND STANDARDS PREPARATION:**

1. **Compound 1** Sample and Working Standard Preparation:

30

Weigh (in duplicate), to the nearest 0.1 mg, about 20 mg of **Compound 1** working standard and samples and add to individual 100 mL flasks. Duplicate weights should be prepared for the working standard and each sample lot. Pipet 10 mL of THF to each flask, sonicate for about 1 min, add enough mobile phase to fill each flask approximately 80 percent of capacity, shake, and allow to equilibrate to room

temperature. Dilute to volume (QS) with mobile phase. (See operator's note #1). Designate the working standard solution as **POT1**. Designate the sample solution as **A1**.

2. Preparation of Working Standard for **Compound 6**

5

	<u>COMPOUND</u>	<u>CODE</u>	<u>SAMPLE WEIGHT</u>	<u>FLASK</u>
10	Compound 6	Ox	10 mg	100mL

15

Pipet 10 mL of THF to the flask, sonicate for 1 minute, add enough mobile phase to fill the flask to approximately 80 percent of capacity, shake, and allow to warm to room temperature. Dilute to volume with additional mobile phase and label as **PUR 1**.

B. Further dilute the **PUR1** solution to yield **PUR2** solutions as follows:  
Label the flask with as solution **PUR2**.

20

	<u>COMPOUND</u>	<u>CODE</u>	<u>TRANSFER VOLUME</u>	<u>FLASK SIZE</u>
	Compound 6	Ox	2 mL <b>PUR1</b>	100 mL

25

Dilute the solution (**PUR2**) flask to volume with mobile phase.  
C. Prepare the final concentration of the oxindole impurity by diluting the **PUR2** solution to yield a **PUR3** solution:

	<u>COMPOUND</u>	<u>CODE</u>	<u>TRANSFER VOLUME</u>	<u>FLASK SIZE</u>
30	Compound 6	Ox	2 mL <b>PUR2</b>	100mL

Pipet the indicated volume of **PUR2** solution into the flask and add 10 mL of THF. Fill each flask to approximately 80 percent of capacity with mobile phase, and

allow to warm to room temperature. Dilute to volume with mobile phase. Label the flask as solution **PUR3**.

**SYSTEM SUITABILITY:**

- 5 A complete system suitability should be determined prior to the initial assay and after any significant change to the system. For these criteria, see the **SYSTEM SUITABILITY** section that follows in this procedure.

**SYSTEM SUITABILITY PRIOR TO EACH ANALYSIS:**

The following criteria must be achieved prior to every analysis made with this procedure.

- 10 1. Calculate the resolution between **Compound 1** and **Compound 6** by using the preparation specified in the **SYSTEM SUITABILITY** section.
2. Verify that an adequate limit of quantitation (LOQ) is achieved. Using the solution **PUR3**, perform 2 replicate injections. The areas of the **Compound 6** peaks should agree within 25%. Calculate the % area agreement as follows:

15

$$\% \text{ area agreement} = \frac{2(A-B) \times 100}{A+B}$$

Where: A = Area of oxindole (**Compound 6**) peak in injection 1

B = Area of oxindole (**Compound 6**) peak in injection 2

- 20 3. For the working standard solution of **Compound 1 POT1** measure the retention time for **Compound 1**. The retention time should be within the range of 17-21 minutes.

**PROCEDURE:**

- 25 1. Inject purity standard **PUR3**, four samples (A), **PUR3**, four samples, etc. No more than four samples may be injected between standards. For the purity standard (**PUR3**), measure the peak area and retention time of each specified impurity. For each sample injection, measure the retention time and the chromatographic peak area of each peak observed (See operator's note #3).

**IDENTITY TEST:**

- 30 This test is satisfactorily met if the **Compound 1** sample solution (A) under test exhibits a major peak whose retention time is identical ( $\pm 2\%$ ) to that of the **Compound 1** working standard solution (**POT1**).

CALCULATIONS:PURITY:

1. For each sample, establish the presence of oxindole, if any, according to the relative retention time indicated in the table found in the CHROMATOGRAPHIC CONDITIONS section, and by comparison to the retention time of the peaks in the respective purity standard (PUR3). Their identities are established if the retention time does not differ by more than 2% (sample peak versus peak in the specified impurity standard). Specified impurities are to be quantified with the purity standard (PUR3).

2. Calculate the standard response factor for oxindole:

$$SR_i = (A_i)(DF)/(W_i)(PF_i)$$

where:  $SR_i$  = Standard response factor of oxindole  
 $A_i$  = Area of the specified impurity in PUR3  
 $W_i$  = Weight (mg) of the oxindole  
 $PF_i$  = Potency Factor of the oxindole working standard (e.g., 0.993).  
 $DF$  = Dilution factors for oxindole:  
 = 250,000

3. Calculate the percent of oxindole in the following manner:

$$\% \text{ oxindole} = (A(s))(DF)(100)/(SR_i(\text{avg}))(W_s)$$

where:  $SR_i(\text{avg})$  = Average standard response for the oxindole working standard.  
 $W_s$  = Weight of the oxindole sample in mg  
 $A(s)$  = Area of the oxindole in the sample.  
 $DF$  = Dilution factor = 100  
 100 = Conversion to %

30 System Suitability:

The criteria set forth below establish chromatographic conditions that ensure the system is operating in a manner suitable to carry out the procedure. If any of these are failed, appropriate adjustments to the system should be made prior to proceeding. System suitability should be assessed prior to the first analysis on the



system or after any significant change (e.g., replacement of column, repair of autosampler, etc.).

## 1. Reproducibility

5 Perform five replicate injections of the **Compound 1** working standard solution. Measure the peak area for **Compound 1**. The relative standard deviation (coefficient of variation) of the peak areas for **Compound 1** should not exceed 2.0%.

10 Perform six replicate injections of the purity standard solution (PUR3) containing  
Compound 6. Measure the peak area for Compound 6. The relative standard  
deviation (coefficient of variation) of the peak areas for Compound 6 should not  
exceed 15%.

15      2.      **Efficiency**

Calculate the number of theoretical plates (N) for the chromatographic column using one representative injection of the **Compound 1** working standard solution. The number of theoretical plates should be equal to or greater than 6,000.

20  $N = 16 (t/W)^2$

### 3. Peak Asymmetry

25 Calculate the peak asymmetry (T) for the **Compound 1** peak using one representative injection of the working standard solution. The peak asymmetry should not be more than 2.0.

$$T = (W_{0.05} / 2f)$$

30      4.      Resolution

Calculate the resolution (R) between **Compound 1** and **Compound 6** by preparing a sample of **Compound 1** spiked with 0.1% (w/w) of **Compound 6** as follows:

Prepare a solution of **Compound 1** as directed above under Step #1 of the SAMPLE AND STANDARDS PREPARATION. Before diluting to volume, add 10 mL of the **PUR2** solution prepared in Step B in the preparation of impurities standards section above.

5

The resolution between **Compound 1** and **Compound 6** should be  $> 1.0$ .

$$R = (2(t_2 - t_1))/(W_2 + W_1)$$

5. The definitions of the terms used in this section are:

10

$t$  = Retention time measured from time of injection to time of elution of peak maximum

$W$  = Width of peak measured by extrapolating the relatively straight sides to baseline

15

OPERATOR NOTES:

1. The time required to equilibrate normal phase columns generally is longer than for reversed phase columns. A new column initially should be washed with 4 liters of mobile phase. The last peak pair to be resolved during equilibration are **Compound 6** and **Compound 1**. Make several injections of the reference for equilibration.
2. Reference and Sample make-up (diluting and warming to room temperature) should be done at the same time to insure that volumes are the same.
3. A THF blank (10 mL THF to volume with mobile phase) may be run to insure that there is no interference with any peaks of interest.

20

25

As described above, this invention also provides an additional method to control for low des-chloro ziprasidone by ensuring low des-chloro Step 2 intermediate (**Compound 8** of Scheme 2, 5-(2-chloroethyl)-oxindole). This is an important control point, due to the opportunity to monitor the process following the reduction of the carbonyl. Reductive conditions also result in hydro-dehalogenation, and thus in synthesis depicted in Scheme 1 loss of the chlorine desired at the C-6 position.

30

Analytical methodology, the HPLC method described above, was developed to evaluate 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one, compound **3** of

Scheme 1, for assay and purity. One important function of this methodology, is to detect ppm levels of compound 8; the des-chloro impurity, in the isolated material. The procedure "Detection Method A" set forth below exemplifies how an analyst might specifically apply this HPLC detection method to quantitate low levels of compound 8 without interference from other process-related impurities. This Example is not intended to and should not be construed to limit the invention described more fully herein and claimed below:

**Detection Method A:**

Following system suitability, the following analytical chromatographic HPLC method can be used to quantify the levels of des-chloro (Compound 8) in Compound 3.

**APPARATUS:**

1. Suitable HPLC, equipped with standard equipment
2. Column heater, capable of operating at 35 °C, e.g., BAS Temperature controller Model LC22A
3. Mobile phase preheater block: e.g., Bioanalytical Systems, Inc. (BAS), Cat # EW8146

Note: This is required to improve column efficiency.

4. Column - Zorbax-SB-CN (Catalog No. 883975.905) 15 cm length x 4.6 mm I.D. (Available from Mac Mod Analytical, Chadds Ford, PA.)

**REAGENTS:**

1. 0.05 M Monobasic Potassium Phosphate ( $\text{KH}_2\text{PO}_4$ ), pH = 6.0 Buffer Solution  
Dissolve 6.8 g  $\text{KH}_2\text{PO}_4$  in one liter of purified water. Adjust the pH of solution to  $6.0 \pm 0.1$  with 5 N potassium hydroxide solution. Larger volumes may be prepared as needed.
2. Mobile Phase: (75 : 15 : 10 v/v/v) 0.05 M  $\text{KH}_2\text{PO}_4$ , pH= 6.0 : Acetonitrile:Methanol

Filter and degas under reduced pressure with stirring or ultrasonic agitation for about 5 minutes. Larger volumes of mobile phase may be prepared using the appropriate amounts of the components.

CHROMATOGRAPHIC CONDITIONS:

	<u>Parameter</u>	-	<u>Set Point</u>	<u>Variation</u>
5	Mobile Phase	-	as above	± 2% ACN and MeOH
	Column Temperature	-	35 °C	±5 °C
	Detection	-	UV, 254nm	+ 5%
	Flow Rate	-	1.0 mL/min	±0.1 mL/min
	Injection Volume	-	20 µL	permissible
10	Quantification Method	-	Area	-
	Run Time	-	60 minutes	Approximately

Under the above-cited conditions Compound 8 will elute in approximately 8-10 minutes. Relative retention times of specified impurities are tabulated below.

	<u>Compound</u>	<u>Rr</u>
	Compound 1	0.36
5	Compound 2	0.45
	Compound 8	0.49
	Compound 3	1.00

**Note:** Relative Retention Time (Rr) = retention time of specified impurity peak  
10 relative to retention time of Compound 3

**PREPARATION OF REFERENCE STANDARD SOLUTIONS:**

**Compound 8 Standard:** Compound 8 standard into a 200 mL volumetric flask.  
Add approximately 20 mL THF and sonicate until sample has completely dissolved  
15 (approx. 1 min.). Dilute to volume with methanol. Mix well with additional shaking and  
inversions. Identify this as Compound 8 solution F1. The concentration of Compound  
8 in solution F1 is about 0.1 mg/mL.

Dilute 2 mL of F1 to 100 mL with methanol and mix well. Identify this as  
Compound 8 solution F2.

20 The concentration of Compound 8 in solution F2 is about 0.002 mg/mL.

**PREPARATION OF SAMPLE SOLUTIONS:**

**Compound 3 Sample (for Compound 8 determination):**

Prepare one test solution per sample. Weigh approximately 50 mg (record to  
25 the nearest 0.1 mg) of the Compound 3 sample into a 50 mL volumetric flask. Add  
approximately 20 mL of THF and sonicate until the sample has completely dissolved  
(approx. 2 min.). Dilute to volume with mobile phase. Mix well with additional shaking  
and inversions. Identify this as Compound 3, Solution I (Sample Solution I). The  
concentration of solution I (Sample Solution I) is about 1.0 mg/mL.

30 **Note:** Solution I is stable for up to 24 hours under normal laboratory conditions.

**SYSTEM SUITABILITY:**

A complete system suitability should be determined prior to the initial assay and after  
any significant change to the system. For these criteria, see the SYSTEM SUITABILITY  
section at the end of this procedure.

SYSTEM SUITABILITY CHECK PRIOR TO EACH ANALYSIS:

The following criteria must be achieved prior to every analysis made with this procedure.

5

1. Calculate the resolution (R ) between COMPOUND 2 and COMPOUND 8 using the **PUR1** standard. The resolution between this pair of peaks should be  $\geq 1.0$ .

2. Verify that an adequate limit of quantitation (LOQ) is achieved. Using solution **PUR1**, perform 2 replicate injections. The COMPOUND 8 peak areas should agree within  $\pm 20\%$ . Calculate the % area agreement as follows:

10

$$\% \text{ area agreement} = \frac{2(A-B) \times 100}{A+B}$$

15

A = COMPOUND 8 peak area from the 1st injection.

B = COMPOUND 8 peak area from the 2nd injection.

3. For the standard solution of COMPOUND 3 **A1**, measure the retention time for COMPOUND 3. The retention time should be within the range of 16 - 24 minutes.

20

PROCEDURE:

Determine HPLC system suitability using the guidelines and procedures for running system suitability presented later in this test procedure. LOQ, resolution and retention time tests, as described in the **SYSTEM SUITABILITY CHECK PRIOR TO EACH ANALYSIS** sections, must be performed each time the system is used. The remaining criteria for system suitability should be assessed prior to the first analysis on the system and after any significant change

25

EVALUATION OF COMPOUND 8 CONTENT IN COMPOUND 3 SAMPLE:

(Solution (I))

30

Inject 20  $\mu\text{L}$  aliquots of the sample solution (I). Measure the area of the chromatographic peak of Compound 8 from each injection. Determine the Compound 8 levels present in the test sample (I) as described in the **CALCULATIONS** section.

**CALCULATIONS:****Calculation of COMPOUND 8 Content:**

1. Determine the response factor for COMPOUND 8 as follows:

$$R_{pur1} = \frac{A_{pur1} \times DF}{W_{pur1} \times PF}$$

where:  $A_{pur1}$  = Peak area of the impurity (COMPOUND 8) in PUR1

$W_{pur1}$  = Weight of the impurity (COMPOUND 8) in PUR1

PF = Potency factor for COMPOUND 8 or COMPOUND 2 (e.g., 0.993)

DF = Dilution Factor -

COMPOUND 8  $\Rightarrow 2 \times 10^5$

2. Determine COMPOUND 8 content as follows:

$$\% \text{ w/w} = \frac{A_c \times 50 \times 100}{R_{pur1} \times W_{s2}}$$

where:  $A_c$  = Peak area of COMPOUND 8 in Sample I

$R_{pur1}$  = Response factor for COMPOUND 8 in standard \*

$W_{s2}$  = Weight of COMPOUND 3 in sample I in mg

50 = Dilution Factor

100 = Conversion to percent

- \* Use the average response factor for all PUR1 injections made throughout the analysis.

**SYSTEM SUITABILITY:**

The criteria set forth below establish chromatographic conditions that ensure the system is operating in a manner suitable to carry out the procedure. If any of these are failed, appropriate adjustments to the system should be made prior to proceeding. System suitability

should be assessed prior to the first analysis on the system or analysis on the system or after any significant change (e.g., replacement of column, repair of autosampler, etc.).

#### 1. Precision of Injection

- 5        Assay: Perform 5 replicate injections of the COMPOUND 3 standard A1. Measure the area of each COMPOUND 3 peak. The relative standard deviation of the peak areas should not exceed 1.0%.

- Purity Evaluation: Perform 6 replicate injections of the PUR1 standard solution. Measure the area of each COMPOUND 8 peak. The relative standard deviation of the peak  
10       area should not exceed 10%.

#### 2. Efficiency

- Calculate the number of theoretical plates (N) for the chromatographic column using  
15       the COMPOUND 3 peak in standard A1. The number of theoretical plates should not be less than 5000 when calculated by the tangent method.

$$N = 16 \left( \frac{t}{W} \right)^2$$

- 20       t = Retention time measured from time of injection to time of elution of peak maximum.

- W = Width of peak measured by extrapolating the relatively straight sides to baseline.

#### 3. Retention Time

- 25       Measure the retention time for the COMPOUND 3 peak in standard A1. The retention time for the COMPOUND 3 peak should be within the range of 16 - 24 minutes.



#### 4. Peak Asymmetry

Calculate the peak asymmetry (T) for the COMPOUND 3 peak in standard A1. The peak asymmetry should not be more than 2.0.

$$T = \frac{W_{0.05}}{2f}$$

5

T = Tailing factor

$W_{0.05}$  = Peak width at 5% height

f = Distance from the peak maximum to the leading edge of the peak, measured at 5% peak height

10

#### 5. Resolution

Calculate the resolution (R) between COMPOUND 2 and COMPOUND 8 in PUR1.

15 The resolution between this pair of peaks should be  $\geq 1.0$ .

$$R = \frac{2(t_2 - t_1)}{W_1 + W_2}$$

t = Peak retention time

20

W = Peak width at baseline (measured by extrapolation to the baseline of tangents to the inflection points, for each of component)

25

As part of the development of this chemistry, multiple alternative synthetic methods were analyzed for the reduction of compound 2 to compound 3 in Scheme 1 above. For example, catalytic hydrogenation of the carbonyl of Compound 2 was tested using 5% palladium on carbon, palladium on alumina, platinum on carbon, or platinum on alumina. The palladium experiments were repeated at 10% catalyst loading, and in each case dehalogenation was a problem. We found that triethylsilane (TES) in the presence of a strong acid gives reduction of the carbonyl, without contaminate hydro de-halogenation. Therefore, this type of reduction is

30

preferred for the production of compound 3. Further, trifluoroacetic acid or methanesulfonic acid have been found to be preferable strong acids for this chemistry, due to avoiding the formation of des-chloro impurities such as Compound 8.

5 A purification was developed to re-process lots which we considered high in the des-chloro analogs. Processes were developed to rework Compounds 1 or 3. We designed and conducted an experimental program testing purification of the starting material, each intermediate and the final drug to determine the most efficient point in the synthesis of ziprasidone to remove des-chloro compounds if present. We  
10 found that it is most efficient to remove the impurities by recrystallization and/or reslurry of Compounds 1 or 3. Recrystallization of Compounds 2, 4, or the final drug were very inefficient, giving very small reductions in the levels of the des-chloro impurities.

15 Specifically, many conditions were tested to purify Compound 3 including, acetonitrile, methylene chloride/toluene, ethyl acetate/hexanes, isopropyl alcohol, toluene, THF, isopropyl alcohol/DMAC (dimethylacetamide), methanol, isopropyl alcohol/acetic acid, and acetonitrile/water. The purification experiments to ascertain the preferred recrystallization and reslurry conditions for Compound 3 are summarized in Table 1, below. Overall most of the solvents tested were inefficient at  
20 removing Compound 6 from Compound 3. The recrystallization/reslurry from acetonitrile/water was superior to the other procedures tested, giving a reduction in the level of the impurity from ~1300 ppm to ~ 300 ppm. Thus this recrystallization and/or reslurry is preferred for control of the des-chloro impurity 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one.

**Table 1: Purification of Compound 3**

<u>Conditions</u>	<u>Yield</u>	<u>Color</u>	<u>Compound 8</u> (ppm)
Acetonitrile 10v	92.0%	Off-white	870
CH <sub>2</sub> Cl <sub>2</sub> /Toluene 5:5	94.5%	Off-white	1200
EtOAc Distill add Hexane	98.0%	White	1100
Isopropyl alcohol 20 vol	89.3%	Off-white	770
Toluene 20 vol	96.7%	Off-white	930
CH <sub>3</sub> CN/H <sub>2</sub> O 9:1, ¾ hr	92.7%	Off-white	500
CH <sub>3</sub> CN 112 vol, 5% Darco	90.5%	Off-white	430
THF, Darco KB-B	76.6%	White	780
THF	81.2%	White	720
CH <sub>3</sub> CN/H <sub>2</sub> O 9:1, 4 hrs	94.5%	White	440
IPO/DMAC 7:3	78.5%	White	480
10 vol CH <sub>3</sub> OH	94.8%	White	840
IPO/HOAc 4:2	92.5%	Pink	540
CH <sub>3</sub> CN/H <sub>2</sub> O 8:2, 18 hrs	95.8%	White	240
CH <sub>3</sub> CN/H <sub>2</sub> O 9:1, 18 hrs	94.3%	White	230

The ziprasidone drug substance of this invention may be administered as a neuroleptic agent as indicated herein as described in, for example, United States Patent 4,831,031, *supra*. Administration to a mammalian subject, including a human, may be alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents in a pharmaceutical composition, in accordance with standard pharmaceutical practice. The pharmaceutical compositions may be administered orally or parenterally including intravenously or intramuscularly. Suitable pharmaceutical carriers include solid diluents or fillers, and sterile aqueous solutions and various organic solvents. The pharmaceutical compositions are then readily administered in a variety of dosage forms, such as tablets, powders, lozenges, syrups, and injectable solutions. These pharmaceutical compositions, if desired, may contain additional ingredients such as flavorings, binders and excipients. Thus, for purposes of oral administration, tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrants such as starch, alginic acid and certain complex silicates, together with binding agents such as

polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid materials of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Preferred materials for this include lactose or milk sugar and high molecular weight  
5 polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration, the ziprasidone drug substance therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if desired, emulsifying or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin and combinations thereof.

For parenteral administration, solution or suspension of the ziprasidone drug substance  
10 in sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solution may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by  
15 standard techniques known to those skilled in the art.

The effective dosage of ziprasidone depends on the intended route of administration and other factors such as the indication being treated and the age and weight of the subject, as generally known. In general, a daily dosage will be in the range of from about 0.5mg of ziprasidone drug substance to about 500 mg, in single or divided doses, preferably from about  
20 10 mg to about 200 mg per day. Presently, Geodon<sup>TM</sup> is approved in the United States for schizophrenia treatment in capsule form for oral administration comprising the ziprasidone hydrochloride monohydrate form of ziprasidone. The capsules are available in 20, 40, 60, and 80 mg of ziprasidone drug substance dosage forms. A typical daily dose for schizophrenia treatment based on a weight of about 70kg for a patient is preferably from about 20 mg twice  
25 per day to about 100 mg ziprasidone drug substance twice per day, more preferably about 20 mg twice per day to about 80 mg twice per day. However, it is appreciated that the dose and dosing regimen of ziprasidone drug substance may be varied from the aforementioned ranges and regimens by a physician of ordinary skill in the art, depending on the particular circumstances of any specific patient.

30 The following Examples illustrate the present invention. It is to be understood, however, that the invention, as fully described herein and as recited in the claims, is not intended to be limited by the details of the following Examples.

### EXAMPLES

#### Example 1: Synthesis of Ziprasidone

##### Step 1: Friedel-Crafts Acylation of 6-chloro-1,3-dihydro-2H-indol-2-one

Methylene chloride (310 L) and aluminum chloride (172.3 kg) were combined. Chloroacetyl chloride (66.7 kg) was added, and the resulting mixture was stirred for 45 minutes. 6-Chloro-1,3-dihydro-2H-indol-2-one (61.8 kg) was added. The reaction mixture was stirred at 28 to 32° C for 19.5 hours and then cooled to 15 to 20° C. Water (805 L) was cooled to 5 to 10° C. The reaction was quenched by the slow addition of the reaction mixture to the cold water. After the quench was complete, the mixture was heated to reflux, and the methylene chloride was removed by atmospheric distillation at 43 to 57° C. The resulting mixture was cooled to 15 to 20° C and stirred for 1 hour. The solids were isolated by filtration and washed with water (114 L) followed by methanol (114 L). The solids were dried in a suitable dryer.

6-Chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one, yield: 91.3 kg (101.4%). Note: A weight yield in excess of 100% resulted due to small amounts of residual salts which were removed in the following step.

The resulting 6-chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one was carried through the following step in portions, one of which is detailed below.

##### Step 2: Trifluoroacetic Acid/Silane Reduction of 6-Chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one

Trifluoroacetic acid (278 kg) and (74.2 kg) were combined and stirred slowly at 24 to 28° C. Triethylsilane (77.9 kg) was charged to the stirring mixture. The reaction temperature was allowed to exotherm slightly during this addition and was maintained between 50 to 62° C during the reaction period. The reaction mixture was stirred for 8 hours, cooled to 38° C, and sampled for reaction completion. The reaction mixture was stirred at 50 to 54° C for an additional 3 hours. After the reaction was determined to be complete, the reaction mixture was cooled to 18° C, and quenched with water (594 L). The resulting slurry was stirred for 30 minutes at 10 to 15° C, and the solids were isolated by filtration. The product was rinsed from the tank and the product cake was washed with water (83 L) followed by methanol (76 L).

In each of two batches of equal size, tetrahydrofuran (742 L), Darco KB-B (1.9 kg), and the wet product cake were combined and heated to reflux. The resulting

mixture was stirred at reflux for 30 minutes and filtered through a sparkler filter (pre-coated with filteraid) at 50 to 60° C to remove the carbon. The tank and sparkler were rinsed with hot tetrahydrofuran (38 L). Following the filtration the two batches were combined. The solution was concentrated *in vacuo* and stirred at 4 to 5° C for 1 hour. The solids were isolated by filtration and washed with cold tetrahydrofuran (38 L). The solids were dried *in vacuo* at 45 to 73° C until a loss on drying of 0.45% was achieved, giving 6-Chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one, yield: 60.1 kg (85.9%).

The resulting 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one was combined with material of comparable quality and carried through the following step.

Step 3: Coupling of 6-Chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one and 3-(1-Piperazinyl)-1,2-benzisothiazole Monohydrochloride

Water (780 L) and sodium carbonate (126.0 kg) were combined and the mixture was stirred to dissolve. 3-(1-Piperazinyl)-1,2-benzisothiazole monohydrochloride (155.0 kg) and 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one (150.4 kg) were added, and the reaction mixture was heated to reflux (~100° C). After 24 and 28 hours, the reaction slurry was sampled for reaction completion assay. The reaction was determined to be complete after the assay of the second sample. Water (1251 L) was added and the slurry was cooled to temperatures between 18 to 22° C. The solids were isolated by filtration and washed with water (302 L). The water wet solids were combined with isopropanol (940 L) and the resulting mixture was stirred for approximately 2 hours at ambient temperature. The solids were isolated by filtration, washed with isopropanol (89 L), and dried *in vacuo* at less than 43° C, giving 5-[2-[4-(2,3-Benzisothiazol-3-yl)-1-piperazinyl]ethyl]-6-chloro-1,3-dihydro-2*H*-indol-2-one, yield: 202.8 kg (80.8%).

The resulting 5-[2-[4-(2,3-benzisothiazol-3-yl)-1-piperazinyl]ethyl]-6-chloro-1,3-dihydro-2*H*-indol-2-one was divided into two portions. These batches were carried separately through the following additional purification and resulted in material of comparable quality. The processing of one of these batches is detailed below.

Step 3R: Purification of 5-[2-[4-(2,3-Benzisothiazol-3-yl)-1-piperazinyl]ethyl]-6-chloro-1,3-dihydro-2*H*-indol-2-one

5- $[2-[4-(2,3\text{-Benzisothiazol-3-yl})-1\text{-piperazinyl}]ethyl]-6\text{-chloro-1,3-dihydro-2H-indol-2-one}$  (51 kg), filteraid (4 kg) and tetrahydrofuran (2678 L) were combined. The mixture was heated to reflux ( $\sim 65^\circ\text{C}$ ) for  $\sim 1$  hour, filtered while maintaining the temperature above  $55^\circ\text{C}$ , and rinsed with tetrahydrofuran (570 L). The product rich filtrate was partially concentrated *in vacuo*. 5- $[2-[4-(2,3\text{-Benzisothiazol-3-yl})-1\text{-piperazinyl}]ethyl]-6\text{-chloro-1,3-dihydro-2H-indol-2-one}$  (51 kg), filteraid (4 kg) and tetrahydrofuran (2675 L) were combined. The mixture was heated to reflux ( $\sim 65^\circ\text{C}$ ) for  $\sim 1$  hour, filtered while maintaining the temperature above  $55^\circ\text{C}$ , and rinsed with tetrahydrofuran (560 L). The product rich filtrate was combined with the partially concentrated mixture above and concentrated *in vacuo*. The resulting mixture was cooled to 0 to  $5^\circ\text{C}$ . The solids were isolated by filtration, washed with filtered tetrahydrofuran (113 L), and dried *in vacuo* at less than  $41^\circ\text{C}$ , giving ziprasidone, free base, yield: 79.3 kg (77.7 %).

A portion of the batch was combined with material of comparable quality which had been recrystallized separately and the batch was carried through the following step.

**Example 2: Crystallization Salt Formation of Ziprasidone Hydrochloride Monohydrate**

Tetrahydrofuran (2715 L), water (307 L), and 5- $[2-[4-(2,3\text{-benzisothiazol-3-yl})-1\text{-piperazinyl}]ethyl]-6\text{-chloro-1,3-dihydro-2H-indol-2-one}$  (100.0 kg) were combined, heated to reflux ( $\sim 64^\circ\text{C}$ ), and stirred for  $\sim 30$  minutes. The solution was filtered and rinsed with tetrahydrofuran (358 L).

Water (203 L) and concentrated hydrochloric acid (29 L) were combined and stirred at ambient temperature. The resulting aqueous hydrochloric acid solution was charged to the 5- $[2-[4-(2,3\text{-benzisothiazol-3-yl})-1\text{-piperazinyl}]ethyl]-6\text{-chloro-1,3-dihydro-2H-indol-2-one}$  solution over a period of 27 minutes. The reaction mixture was cooled to temperatures between 1 and  $5^\circ\text{C}$  over a period of  $\sim 2$  hours. The mixture was stirred between 1 and  $5^\circ\text{C}$  for  $\sim 10$  hours. The solids were isolated by filtration, washed with cold tetrahydrofuran (358 L), and dried until a water content of 4.1% was obtained.

Ziprasidone Hydrochloride Monohydrate, yield: 108.6 kg (96.0 % weight yield)

The solids were milled on a Bauermeister mill.

**Example 3: Purification of 6-Chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one To Remove 5-(2-Chloroethyl)-1,3-dihydro-2H-indol-2-one**

5 A 100 mL round bottom flask equipped with a magnet stirrer and reflux condenser was charged with 4.0 g (17.4 mmol) of 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one (Compound 3) and 36 mL of acetonitrile and 4.0 mL of water were added. The slurry was gently heated and stirred overnight (~18 hrs at ~78° C). The heating was then removed and the slurry cooled to 0 to 5° C, and stirred for an additional hour. The product was collected by filtration, washed with a small portion  
10 of acetonitrile and the product dried under vacuum at 50° C, to give 3.77 g (94.3% yield) of 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one. The level of the des-chloro impurity had been reduced from 1280 ppm to 230 ppm.

**Example 4: Experimental Determination of Purge Factor for Compound 6 (1,3-Dihydro-2H-indol-2-one)**

15 A batch of 6-chloro-1,3-dihydro-2H-indol-2-one which contained a very high content of 1,3-dihydro-2H-indol-2-one was selected. This was intentionally selected so that higher levels of the impurity would be easier to measure, and to determine the purge factor for this impurity. An additional reason for this strategy of starting with material which was very high in the impurity for purposes of determining the purge factor of the impurity was to avoid having the material purge to less than the limit of analytical detection during the synthesis; thus resulting in a zero value in the final product. Since the purge factor is a ratio, it is not meaningful to divide by a zero result. (The material with the high level of impurity was used for this experiment but  
20 was NOT subsequently used in any studies with human subjects.) A batch of 6-chloro-1,3-dihydro-2H-indol-2-one which contained 4000 ppm of 1,3-dihydro-2H-indol-2-one was processed through the standard synthetic process according to Examples 1 and 2 above.

25 Following the first two steps of the synthesis, the level of the corresponding des-chloro impurity was measured, using the method described. It was found that 1700 ppm of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one (Compound 8 of Scheme 2, above) was present in 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one (Compound 3 of Scheme 1, above). The processing was continued to 5-[4-(1,2)-benzisothiazol-3-yl]-1-piperazinylethyl]-6-chloro-1,3-dihydro-2H-indol-2-one  
30



hydrochloride monohydrate, where it was determined that 600 ppm of 5-[2-[4-(1,2)-benzothiazol-3-yl]-1-piperazinyl]ethyl]-1,3-dihydro-2H-indol-2-one (Compound 9 of Scheme 2, above) was present.

Thus the purge factor through the entire synthesis for the des-chloro analogs was from 4000 ppm to 600 ppm, or approximately a 6-fold decrease. Minor run to run variations in processing can lead to small differences in the yield and quality of the materials produced. A 20% error in the reproducibility of the impurity formation, that is if 500 ppm in one run expecting between 400 and 600 ppm in other experiments, is then allowed for. In the case of the synthesis described in Examples 1 and 2, with 5 processing steps, the additive experimental error could result in as much as a 2-fold difference in the level of the impurity. Thus, for the purpose of setting the upper limit, where the drug is going to be used by human subjects a conservative 3-fold purge factor was utilized. Therefore, to insure that the product produced would not contain over 100 ppm of 5-[2-[4-(1,2)-benzothiazol-3-yl]-1-piperazinyl]ethyl]-1,3-dihydro-2H-indol-2-one (Compound 9), a limit of 300 ppm of 1,3-dihydro-2H-indol-2-one (Compound 6) in 6-chloro-1,3-dihydro-2H-indol-2-one (Compound 1) was determined.

CLAIMS

1. A composition comprising ziprasidone and an amount of des-chloro ziprasidone selected from not greater than about 1000 ppm, not greater than about 500 ppm, and not greater than about 100 ppm.

5 2. A composition according to claim 1, wherein the ziprasidone is ziprasidone free base, ziprasidone hydrochloride monohydrate, ziprasidone mesylate dihydrate, or ziprasidone mesylate trihydrate.

10 3. A pharmaceutical composition for treating in a mammal a disorder or condition selected from schizophrenia, anxiety, migraine pain, Tourette's Syndrome, glaucoma, ischemic retinopathy, dementia of the Alzheimer's type, a bipolar disorder, a mood disorder, agoraphobia, social phobia, panic disorder, post-traumatic stress disorder, acute stress disorder, substance-induced anxiety disorder, an anxiety disorders not otherwise specified (NOS), dyskinesias, a behavioral manifestation of mental retardation, conduct disorder, and autistic disorder comprising an amount of the composition of claim 1 effective in treating said disorder or condition and a pharmaceutically acceptable carrier.

15 4. A method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone selected from not greater than about

A) 1000 ppm,

20 B) not greater than about 500 ppm, and

C) not greater than about 100 ppm

which method comprises:

a) obtaining one or more samples of one or more 6-chloro-1,3-dihydro-2H-indol-2-one batches;

25 b) measuring the level of oxindole impurity in each of the samples of (a);

c) selecting a 6-chloro-1,3-dihydro-2H-indol-2-one batch that comprises a level of oxindole of

for (A), not greater than about 0.3% based on the measurement or measurements conducted in (b),

30 for (B), not greater than about 0.15% based on the measurement or measurements conducted in (b), and

for (C), not greater than about 0.03% based on the measurement or measurements conducted in (b); and

d) using the batch selected in (c) to synthesize said ziprasidone composition.

5. A method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

5 a) acylating a composition comprising 6-chloro-1,3-dihydro-2H-indol-2-one and an oxindole impurity with chloroacetyl chloride by Friedel-Crafts Acylation to synthesize a composition comprising 6-chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one;

10 b) treating the composition resulting from (a) to reduce of the oxo of the chloroacetyl group therein to form a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity;

c) isolating a sample of the composition resulting from (b);

d) measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity in the isolated sample from (c);

15 e) determining whether or not the quantity in (d) is not greater than about 0.28%; and

f) purifying by recrystallization and/or reslurry the composition resulting from (b) if the quantity measured in (d) is greater than about 0.28% until the quantity of the 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity is not greater than about 0.28%, and synthesizing a ziprasidone composition from the composition so purified; or,

20 g) if the quantity in (d) is not greater than about 0.28%, synthesizing a ziprasidone composition from the composition of (b).

6. A method using HPLC for measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one, which method comprises

25 a) preparing sample solution from said composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one by dissolving a portion of said composition in an organic solvent, followed by dilution with an organic solvent of the dissolved portion so that a concentration (weight/volume), based on the weight of said portion and volume of solvent, of about 1 mg/mL is obtained;

30 b) running the sample solution through a stable-bond cyano HPLC column using a mobile phase consisting essentially of (75:13-17:8-12 v/v/v) 0.05 M  $\text{KH}_2\text{PO}_4$ , pH=from 5.5-6.5; Acetonitrile:Methanol; at a column temperature of from 30° C to 40° C; with detection by UV light at 254nm UV;

c) detecting a peak appearing at from between 8 to 10 minutes on a chromatogram resulting from (b);

d) measuring the peak area (named  $A_c$ ) of the peak detected in (c);

e) preparing a standard from a composition consisting essentially of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one by dissolving and diluting a portion of said composition in an organic solvent such that the concentration (weight/volume) of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one, based on the weight of the portion and volume of the solvent, is about equal to a selected fraction value at or above which detection of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one is desired;

f) running the standard through a stable-bond cyano HPLC column using a mobile phase consisting essentially of (75:13-17:8-12 v/v/v) 0.05 M  $\text{KH}_2\text{PO}_4$ , pH=from 5.5-6.5, Acetonitrile:Methanol; at a column temperature of from 30° C to 40° C; with detection by UV light at 254nm UV;

g) measuring the peak area (named  $A_{\text{pur1}}$ ) of the peak on a chromatogram resulting from (f); and

h) calculating the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one in the composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one by

i) calculating the Response Factor for 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one according to the following formula:

$$R_{\text{pur1}} = (A_{\text{pur1}})(DF)/(W_{\text{pur1}})(PF)$$

wherein :  $A_{\text{pur1}}$  is as defined above;

$W_{\text{pur1}}$  = weight of composition in the standard;

PF = potency factor for 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one; and

DF = dilution factor for the standard; and

ii) calculating the %w/w of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one according to the following formula:

$$\%w/w = (A_c)(DF)(100)/(R_{\text{pur1}})(W_{S2})$$

wherein :  $A_c$  is as defined above;

$R_{\text{pur1}}$  = Response Factor calculated in (h)(i) above;

$W_{\text{S2}}$  = weight of the portion of the composition used in step (a); and

DF = dilution factor for sample solution.

5           7.       A method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about 1000 ppm, which method comprises:

10               a) reducing a composition comprising 6-chloro-5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one and a 5-(chloroacetyl)-1,3-dihydro-2H-indol-2-one impurity by treatment with triethylsilane in the presence of a strong acid to obtain a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity; and

              b) synthesizing a composition comprising ziprasidone from the composition resulting from (a).

15           8.       A method according to claim 7, further comprising

              i) isolating prior to step (b) a sample of the composition from (a), and measuring the quantity of 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity in said sample;

20               ii) determining whether or not the quantity in (i) is not greater than an amount selected from about 0.28%; and

              iii) purifying by recrystallization and/or reslurry the composition from (a) if the quantity in (i) is greater than about 0.28% until the quantity of the 5-(2-chloroethyl)-1,3-dihydro-2H-indol-2-one impurity is not greater than about 0.28%, and then proceeding to step (b) using the composition from (a) so purified; or

25               iv) if the quantity in (i) is not greater than about 0.28%, then proceeding to step (b).

              9.       A method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than an amount selected from

- 30               A)       about 1000 ppm,  
              B)       about 500 ppm, and  
              C)       about 100 ppm,

which method comprises:

a) purifying a composition comprising 6-chloro-1,3-dihydro-2*H*-indol-2-one and an oxindole impurity until a composition comprising an amount of said oxindole impurity of

for (A), about 0.3%

for (B), about 0.15%,

and for (C), about 0.03%

is obtained; and

b) using the composition resulting from (a) to synthesize a ziprasidone composition.

10. A method of synthesizing a ziprasidone composition that comprises an amount of des-chloro ziprasidone of not greater than about

A) 1000 ppm,

B) 500 ppm, or

C) 100 ppm;

which method comprises:

a) recrystallizing and/or reslurrying a composition comprising 6-chloro-5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one and a 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity until a composition comprising not greater than about

for (A), 0.3%,

for (B), 0.15%, and

for (C), 0.03%

of said 5-(2-chloroethyl)-1,3-dihydro-2*H*-indol-2-one impurity is obtained; and

b) using the composition resulting from (a) to synthesize a ziprasidone composition.

## INTERNATIONAL SEARCH REPORT

Internatic Application No

PCT/IB 03/00642

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K31/496 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ, BIOSIS, EMBASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 338 846 A (BUSCH FRANK R ET AL) 16 August 1994 (1994-08-16) cited in the application the whole document	1-3
X	US 6 110 918 A (BUSCH FRANK R ET AL) 29 August 2000 (2000-08-29) cited in the application the whole document	1-3
X	EP 0 586 191 A (PFIZER) 9 March 1994 (1994-03-09) the whole document	1-3
X	EP 0 931 547 A (PFIZER PROD INC) 28 July 1999 (1999-07-28) the whole document page 12, line 37 - line 39	1-3
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

14 May 2003

Date of mailing of the international search report

22/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Economou, D

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 03/00642

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 584 903 A (PFIZER) 2 March 1994 (1994-03-02) cited in the application the whole document ----	1-3
X	US 5 359 068 A (URBAN FRANK J) 25 October 1994 (1994-10-25) cited in the application column 10, line 63 -column 11, line 45 claim 8 ----	1-3
X	WO 97 42191 A (SHINE RUSSELL JAMES ;PFIZER (US); ROSE CAROL ANNE (US); BUSCH FRAN) 13 November 1997 (1997-11-13) cited in the application the whole document -----	1-3



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 03/00642

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
US 5338846	A	16-08-1994	US	5206366 A		27-04-1993
			US	5312925 A		17-05-1994
			CA	2095587 A1		27-02-1994
			AT	206422 T		15-10-2001
			AU	642836 B1		28-10-1993
			BR	9302065 A		26-07-1994
			CN	1083061 A ,B		02-03-1994
			CZ	9300877 A3		13-04-1994
			DE	69330853 D1		08-11-2001
			DE	69330853 T2		02-05-2002
			DK	584903 T3		03-12-2001
			EG	20214 A		30-11-1997
			EP	1029861 A1		23-08-2000
			EP	0584903 A1		02-03-1994
			ES	2161703 T3		16-12-2001
			FI	932012 A		27-02-1994
			HU	65750 A2		28-07-1994
			IL	105622 A		15-06-1998
			JP	2742370 B2		22-04-1998
			JP	6184143 A		05-07-1994
			KR	123441 B1		24-11-1997
			MX	9302813 A1		28-02-1994
			NO	931656 A		28-02-1994
			NZ	247539 A		26-09-1995
			PL	299002 A1		21-03-1994
			PT	584903 T		28-02-2002
			RU	2061695 C1		10-06-1996
			SI	9300287 A		31-03-1994
			SK	48593 A3		06-04-1994
			ZA	9306225 A		27-02-1995
			AU	657231 B2		02-03-1995
			AU	4600493 A		16-06-1994
			BR	9303014 A		15-03-1994
			CA	2105114 A1		02-03-1994
			CN	1089607 A ,B		20-07-1994
			CZ	9301789 A3		13-04-1994
			DE	9312903 U1		05-01-1994
			EG	20251 A		31-05-1998
			EP	0586191 A1		09-03-1994
			FI	933804 A		02-03-1994
			HU	67023 A2		30-01-1995
			IL	106777 A		15-04-1997
			JP	2742372 B2		22-04-1998
			JP	6157521 A		03-06-1994
			MX	9305277 A1		31-03-1994
			NO	933093 A		02-03-1994
			NZ	248543 A		26-07-1995
			PL	300235 A1		05-04-1994
			PL	174396 B1		31-07-1998
			RU	2081116 C1		10-06-1997
US 6110918	A	29-08-2000	AT	236902 T		15-04-2003
			AU	730856 B2		15-03-2001
			AU	1936897 A		26-11-1997
			BG	63601 B1		28-06-2002
			BG	102893 A		30-09-1999
			BR	9708932 A		03-08-1999
			CA	2252895 C		20-08-2002

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 03/00642

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6110918	A	EA 1180 B1	30-10-2000
		EP 0904273 A1	31-03-1999
		HU 9902808 A2	28-03-2000
		IL 126590 A	25-11-2001
		JP 3102896 B2	23-10-2000
		JP 11509865 T	31-08-1999
		NO 985193 A	06-11-1998
		PL 329880 A1	12-04-1999
		SK 150598 A3	13-03-2000
		AP 838 A	03-05-2000
		CN 1216990 A , B	19-05-1999
		CZ 9803494 A3	15-09-1999
		HR 970235 A1	30-06-1998
		WO 9742190 A1	13-11-1997
		KR 2000010822 A	25-02-2000
		NZ 508303 A	27-07-2001
		TR 9802241 T2	22-02-1999
		TW 427989 B	01-04-2001
		ZA 9703875 A	06-11-1998
EP 0586191	A 09-03-1994	US 5312925 A	17-05-1994
		AU 657231 B2	02-03-1995
		AU 4600493 A	16-06-1994
		BR 9303014 A	15-03-1994
		CA 2095587 A1	27-02-1994
		CA 2105114 A1	02-03-1994
		CN 1089607 A , B	20-07-1994
		CZ 9301789 A3	13-04-1994
		DE 9312903 U1	05-01-1994
		EG 20251 A	31-05-1998
		EP 0586191 A1	09-03-1994
		FI 933804 A	02-03-1994
		HU 67023 A2	30-01-1995
		IL 106777 A	15-04-1997
		JP 2742372 B2	22-04-1998
		JP 6157521 A	03-06-1994
		MX 9305277 A1	31-03-1994
		NO 933093 A	02-03-1994
		NZ 248543 A	26-07-1995
		PL 300235 A1	05-04-1994
		PL 174396 B1	31-07-1998
		RU 2081116 C1	10-06-1997
		TW 422845 B	21-02-2001
		US 5338846 A	16-08-1994
		ZA 9306394 A	28-02-1995
EP 0931547	A 28-07-1999	AT 231394 T	15-02-2003
		AU 739472 B2	11-10-2001
		AU 9717098 A	08-07-1999
		DE 69810889 D1	27-02-2003
		DK 931547 T3	24-02-2003
		EP 0931547 A1	28-07-1999
		HU 9802958 A1	28-03-2000
		JP 11246409 A	14-09-1999
		NZ 333436 A	23-06-2000
		US 6245766 B1	12-06-2001
		ZA 9811573 A	19-06-2000

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 03/00642

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0584903	A	02-03-1994	US 5206366 A	27-04-1993
			AT 206422 T	15-10-2001
			AU 642836 B1	28-10-1993
			BR 9302065 A	26-07-1994
			CA 2095587 A1	27-02-1994
			CN 1083061 A , B	02-03-1994
			CZ 9300877 A3	13-04-1994
			DE 69330853 D1	08-11-2001
			DE 69330853 T2	02-05-2002
			DK 584903 T3	03-12-2001
			EG 20214 A	30-11-1997
			EP 1029861 A1	23-08-2000
			EP 0584903 A1	02-03-1994
			ES 2161703 T3	16-12-2001
			FI 932012 A	27-02-1994
			HU 65750 A2	28-07-1994
			IL 105622 A	15-06-1998
			JP 2742370 B2	22-04-1998
			JP 6184143 A	05-07-1994
			KR 123441 B1	24-11-1997
			MX 9302813 A1	28-02-1994
			NO 931656 A	28-02-1994
			NZ 247539 A	26-09-1995
			PL 299002 A1	21-03-1994
			PT 584903 T	28-02-2002
			RU 2061695 C1	10-06-1996
			SI 9300287 A	31-03-1994
			SK 48593 A3	06-04-1994
			US 5338846 A	16-08-1994
			ZA 9306225 A	27-02-1995
US 5359068	A	25-10-1994	CA 2166203 C	24-04-2001
			EP 0706524 A1	17-04-1996
			FI 956279 A	27-12-1995
			WO 9500510 A1	05-01-1995
			JP 2667987 B2	27-10-1997
			JP 8506354 T	09-07-1996
WO 9742191	A	13-11-1997	AP 765 A	17-09-1999
			AU 731267 B2	29-03-2001
			AU 2174797 A	26-11-1997
			BG 63544 B1	30-04-2002
			BG 102892 A	30-09-1999
			BR 9709889 A	10-08-1999
			CN 1216991 A , B	19-05-1999
			CZ 9803493 A3	15-09-1999
			EA 1190 B1	25-12-2000
			EP 0918772 A1	02-06-1999
			HR 970236 A1	30-06-1998
			WO 9742191 A1	13-11-1997
			IL 126591 A	25-11-2001
			JP 11509867 T	31-08-1999
			KR 2000010824 A	25-02-2000
			NO 985194 A	06-11-1998
			NZ 508304 A	25-05-2001
			PL 329884 A1	12-04-1999
			SK 150898 A3	14-02-2000
			TR 9802240 T2	22-02-1999

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 03/00642

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9742191	A	US 6245765 B1 ZA 9703876 A	12-06-2001 06-11-1998